Interactions of Lithium Atoms in a Resonator-Enhanced Dipole Trap

<u>M. Bartenstein</u>, S. Jochim, G. Hendl, J. Hecker Denschlag, A. Mosk¹, M. Weidemüller² and R. Grimm

Institut für Experimentalphysik, Universität Innsbruck Technikerstrasse 25, 6020, Innsbruck, Austria Tel +43-512-5076306, Fax +43-512-5072921 E-mail: Markus.Bartenstein@uibk.ac.at, Website: http://exphys.uibk.ac.at/ultracold ¹FOM Inst. for Plasma Physics, Rijnhuizen, The Netherlands ²Max-Planck-Institut für Kernphysik, Heidelberg, Germany

In order to study the interaction in an ultracold lithium gas, we have developed a novel dipole trap based on the resonant enhancement of a far detuned standing wave in an optical resonator. Light from an ultrastable 2 Watt Nd:YAG laser is coupled into the resonator and enhanced 130-fold, thus creating a trap depth of approximately 1 mK for the lithium atoms. It enables us to trap roughly 10^5 to 10^6 laser-cooled atoms in about 1000 wells of the standing wave[1].



Figure 1: Schematic of the resonator trap.

Current experimental work is being performed with the fermionic isotope, ⁶Li. The goal is to explore a Feshbach resonance that is predicted at a magnetic field of 800 G[2]. In following experiments, the tunability of the interactions in the vicinity of this resonance will be essential to accelerate thermalization of the trapped gas, which is essential for a variety of cooling techniques[3]. Also, a strongly attractive interaction will be necessary to observe a phase transition to a BCS phase at an experimentally feasible temperature.

The diode laser system used for the magneto-optical trap can be easily adapted to trap the bosonic isotope, ⁷Li. With the bosons we plan to explore combinations with magnetic traps[4], a novel collision-assisted cooling scheme[5], and the production of a stable array of small attractively interacting Bose-Einstein condensates.

- [1] A. Mosk et al., Optics Letters, **26**, No. 23 (2001).
- [2] M. Houbiers et al., PRA 57, R1497 (1998).
- [3] N. Poli et al., PRA(R), **65**, 021401 (2002).
- [4] D. M. Stamper-Kurn et al., Phys. Rev. Lett. 81, 2194 (1998).
- [5] G. Ferrari, Eur. Phys. J. D 13, 67 (2001)