

Interference between different Rubidium Dark States

G. Alzetta¹ L. Botti² and A. Rossi²

¹ *INFN UdR Siena - Department of Physics, University of Pisa, Piazza Torricelli 2, I-56126 Pisa, ITALY*

² *INFN UdR Siena - Department of Physics, University of Siena, Via Banchi di Sotto 55, 53100 Siena, ITALY*

We show an experiment in which we have studied the behavior of two different Dark States [1] selected by two diode lasers, one tuned at the D_1 transition and the other at the D_2 . The Dark States are obtained in cells filled with vapor of Rubidium 85 and both are selected in the same spatial region. The modulation frequency ν corresponds to the hyperfine energy separation between the $F=2$ and the $F=3$ levels of the $5S_{1/2}$ ground state, so a double Λ scheme [2] between the $5S_{1/2}$ and the $5P_{1/2}$ states and between the $5S_{1/2}$ and the $5P_{3/2}$ is reproduced. We developed a method to compare and change the phase difference $\gamma = \Theta_{D_1} - \Theta_{D_2}$ between the optical beats emitted by each laser.

Finally we show the behavior of the dark states when are obtained by alone and together. A strong relation between the value of the phase difference γ and the interference between the two dark state is found out. In particular the dark state D_1 coexists with the D_2 when the phase difference γ is π , while they interfere destructively when the phase difference is zero.

Furthermore, as application of the set up used in the experiment described above, we show a way to modulate the laser beam intensity by interaction with the Rb atoms pumped in a dark state by a second laser.

[1] G. Alzetta, A. Gozzini, L. Moi, and G. Orriols, *Nuovo Cimento* **36B**, 5 (1976); G. Alzetta, L. Moi, and G. Orriols, *Nuovo Cimento* **52B**, 209 (1979)

[2] E. Arimondo in *Progress in Optics XXXV*, ed. by E. Wolf (Elsevier Science, 1996), pag.257