

Mixtures of Degenerate Atomic Gases : Bosonic ${}^7\text{Li}$ and Fermionic ${}^6\text{Li}$

J. Cubizolles, T. Bourdel, G. Ferrari, L. Khaykovich, K. Corwin, F. Schreck and C. Salomon

Laboratoire Kastler Brossel, École Normale Supérieure

24 rue Lhomond 75231 Paris Cedex 05

Tel +33-144322019, Fax +33-144323434

E-mail: Julien.Cubizolles@lkb.ens.fr

We designed a system in which two lithium isotopes, the boson ${}^7\text{Li}$ and the fermion ${}^6\text{Li}$ can be trapped in a Magneto-Optical Trap and cooled in a magnetic Ioffe-Pritchard trap [1]. This provides a physical system with several interesting situations ranging from mixture of interacting degenerate quantum gases with different statistics, Bose-Einstein condensate with attractive or repulsive interactions at will, and possibly to pairing of strongly interacting fermions [2] as for the Bardeen-Cooper-Schrieffer (BCS) transition in superconductors.

Since there are no s-wave collisions between identical fermions at zero temperature due to the Pauli exclusion principle we developed a sympathetic cooling scheme in which the bosons are cooled down using the classical evaporative cooling technique and the fermions thermalize by elastic collisions with the bosons. Using ${}^7\text{Li}$ atoms in a hyperfine state where the scattering length is positive (repulsive interactions) we were able to reach the degeneracy regime with both isotopes [3]. The equilibrium temperature of the mixture is $0.28\mu\text{K} \simeq 0.2(1)T_C = 0.2(1)T_F$ where T_C is the BEC transition temperature and T_F the Fermi temperature. The signature of the BEC phase transition is straightforward in the momentum distribution of the bosons whereas precise study of the fermions distribution is needed to observe the effect of Fermi pressure. This mixture offers interesting prospects for the study of phase separation effects [4, 5].




Figure 1: Absorption images of a soliton travelling in the horizontal 1D optical waveguide.

The scattering length can also be tuned by means of a bias magnetic field when a *Feshbach resonance* [6] occurs between two hyperfine states of one or two isotopes. Using such a resonance we could reach Bose-Einstein Condensation with ${}^7\text{Li}$ in an optical trap with a positive scattering length ($a = 2.1$ nm at 665 G) and then tune the scattering length to a negative value ($a = -0.21$ nm at 425 G) [8]. In a one-dimensional configuration, this attractive condensate exhibits a so-called solitonic behaviour. The non-linearity arising from the attractive interaction term compensates the dispersion due to the kinetic energy term resulting in a propagation without spreading [7]. After releasing our attractive condensate in a 1-D optical dipole trap we observed a propagation without spread over a macroscopic distance of 1.1 mm.

- [1] F. Schreck, G. Ferrari, K. L. Corwin, J. Cubizolles, L. Khaykovich, M.-O. Mewes, and C. Salomon, *Phys. Rev. A* **64** 011402 (2001).
- [2] H.T.C. Stoof and M. Houbiers in *Bose-Einstein Condensation in Atomic Gases*, Proceedings of the International School of Physics “Enrico Fermi”, edited by M. Inguscio, S. Stringari and C.E. Wieman (IOS Press, Amsterdam, 1999).
- [3] F. Schreck, L. Khaykovich, K.L. Corwin, G. Ferrari, T. Bourdel, J. Cubizolles and C. Salomon *Phys. Rev. Lett.* **87** 080403 (2001).
- [4] K. Moelmer, *Phys. Rev. Lett.* **80** 1804 (1998).
- [5] Z. Akdeniz, P. Vignolo, A. Minguzzi, and M.P. Tosi cond-mat/0112183.
- [6] A.J. Moerdijk, B.J. Verhaar, and A. Axelsson, *Phys. Rev. A.* **51** 4852 95, Williams, private communications.
- [7] Yvan Castin in *Coherent Atomic Matter Waves*, edited by R. Kaiser, C. Westbrook and F. David (EDP Sciences: Springer-Ferlag 2001).
- [8] L. Khaykovich, F. Schreck, G. Ferrari, T. Bourdel, J. Cubizolles, L. D. Carr, Y. Castin, and C. Salomon, *submitted to Science*.